



On Dividing a Square Into Triangles

Author(s): Paul Monsky

Source: *The American Mathematical Monthly*, Vol. 77, No. 2, (Feb., 1970), pp. 161-164

Published by: Mathematical Association of America

Stable URL: <http://www.jstor.org/stable/2317329>

Accessed: 27/05/2008 23:05

---

Your use of the JSTOR archive indicates your acceptance of JSTOR's Terms and Conditions of Use, available at <http://www.jstor.org/page/info/about/policies/terms.jsp>. JSTOR's Terms and Conditions of Use provides, in part, that unless you have obtained prior permission, you may not download an entire issue of a journal or multiple copies of articles, and you may use content in the JSTOR archive only for your personal, non-commercial use.

Please contact the publisher regarding any further use of this work. Publisher contact information may be obtained at <http://www.jstor.org/action/showPublisher?publisherCode=maa>.

Each copy of any part of a JSTOR transmission must contain the same copyright notice that appears on the screen or printed page of such transmission.

---

JSTOR is a not-for-profit organization founded in 1995 to build trusted digital archives for scholarship. We enable the scholarly community to preserve their work and the materials they rely upon, and to build a common research platform that promotes the discovery and use of these resources. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

countable or locally compact space must be a  $k$ -space in this sense (compare [4], p. 231, Theorem 13); a space in which each compact subset is locally compact is a  $k$ -space if and only if it is a quotient of a locally compact space (compare [1], Theorem 11.9.4); a quotient of any  $k$ -space is a  $k$ -space.

**10. Specialize when necessary.** In some situations  $T_2$  may be needed or convenient. In such cases one specializes; but this is no reason to use  $T_2$  as a blanket assumption from the beginning, any more than prospective use of the Čech compactification justifies dealing only with completely regular spaces from the beginning.

**11. Examples.** Some examples of useful non-Hausdorff topologies are: the topology induced by a non-separating family of seminorms on a vector space, the one point compactification (see [9]), the Zariski topology used by algebraic geometers.

#### References

1. J. Dugundji, *Topology*, Allyn and Bacon, Boston, 1966.
2. L. Gillman and M. Jerison, *Rings of Continuous Functions*, Van Nostrand, Princeton, N. J., 1960.
3. H. Herrlich, On intersections of compact sets, *Duke Math. J.*, 35 (1968) 439–440.
4. J. L. Kelley, *General Topology*, Van Nostrand Princeton, N. J., 1955.
5. J. L. Kelley, I. Namioka *et al.*, *Linear Topological spaces*, Van Nostrand, Princeton, N. J., 1963.
6. N. Levine, When are compact and closed equivalent? *this MONTHLY*, 72 (1965) 41–44.
7. N. Steenrod, A convenient category of topological spaces, *Michigan Math. J.*, 14 (1967) 133–152.
8. A. Wilansky, *Functional Analysis*, Blaisdell, Waltham, Mass., 1964.
9. ———, Between  $T_1$  and  $T_2$ , *this MONTHLY*, 74 (1967) 261–266.

---

## MATHEMATICAL NOTES

EDITED BY DAVID DRASIN

*Manuscripts for this Department should be sent to David Drasin, Division of Mathematical Sciences, Purdue University, Lafayette, IN 47907.*

### ON DIVIDING A SQUARE INTO TRIANGLES

PAUL MONSKY, Brandeis University and Kyoto University

Sometime ago in this *MONTHLY*, Fred Richman and John Thomas [1] asked the following puzzling question:

*Can a square  $S$  be divided into an odd number of nonoverlapping triangles  $T_i$ , all of the same area?*

In [2], the answer was shown to be no, provided  $S = [0, 1] \times [0, 1]$  and the coordinates of the vertices of the  $T_i$  are rational numbers with odd denominators. In this note we shall show that the answer is always no. In fact we shall prove the following more general result.

Suppose that  $S = [0, 1] \times [0, 1]$  is divided into  $m$  nonoverlapping triangles  $T_i$ ; let  $a_i = \text{area } T_i$ . Then there is a polynomial  $f$  with integer co-efficients such that  $f(a_1, \dots, a_m) = 1/2$ .

There are two parts to the proof: one combinatorial, the other valuation theoretic. The combinatorial argument generalizes an argument made in [2]. By itself it may be made to prove the desired result when the vertices of the  $T_i$  all have rational coordinates. But to handle the case of arbitrary vertices, it becomes necessary to argue with "congruences mod 2 in the reals." This is where valuation theory comes in; we make use of absolute values on the reals extending the 2-adic absolute value of the rationals. In the course of the proof the theorem of the extension of valuations plays a remarkable and unexpected role.

We begin with the combinatorial argument. Let  $R$  be a region in the plane bounded by a simple closed polygon. Suppose  $R$  is divided into  $m$  nonoverlapping triangles  $T_i$ . By a *vertex* we shall mean a vertex of some  $T_i$ , by a *face* a face of some  $T_i$  or of  $R$ . Two vertices are called *adjacent* if they are in the same face and the line segment joining them contains no other vertices. A *basic segment* is a line segment joining two adjacent vertices. Note that the boundary of each  $T_i$  is a union of nonoverlapping basic segments; the same is true of the boundary of  $R$ . Suppose now that the vertices are divided into three disjoint sets,  $\mathcal{A}$ ,  $\mathcal{B}$ , and  $\mathcal{C}$ . We shall say that a face or a basic segment is of *type*  $\mathcal{A}\mathcal{B}$  if it has one endpoint in  $\mathcal{A}$  and one in  $\mathcal{B}$ .

LEMMA. *Suppose that no face contains vertices of all three types and that  $R$  has an odd number of faces of type  $\mathcal{A}\mathcal{B}$ . Then some  $T_i$  has vertices of all three types.*

To prove the lemma note the following. A face of type  $\mathcal{A}\mathcal{B}$  contains an odd number of basic segments of type  $\mathcal{A}\mathcal{B}$ , while a face not of type  $\mathcal{A}\mathcal{B}$  contains an even number of basic segments of type  $\mathcal{A}\mathcal{B}$ . (Use the fact that a face contains vertices of at most two types to prove this.) Suppose that no  $T_i$  has vertices of all three types. Then each  $T_i$  has either 0 or 2 faces of type  $\mathcal{A}\mathcal{B}$ . Hence the boundary of  $T_i$  contains an even number of basic segments of type  $\mathcal{A}\mathcal{B}$ . Similarly, the boundary of  $R$  contains an odd number of basic segments of type  $\mathcal{A}\mathcal{B}$ . But this is impossible; in an obvious sense the boundary of  $R$  is congruent to the sum of the boundaries of the  $T_i$  modulo 2.

We now come to the valuation theoretic part of the proof and need to introduce some further terminology. Let  $K$  be a field. By an *ultranorm* (sometimes called a *non-Archimidean absolute value*) on  $K$  we mean a function  $\| \cdot \|$  from  $K$  to the nonnegative real numbers satisfying:

- (1)  $\|xy\| = \|x\| \cdot \|y\|$
- (2)  $\|x + y\| \leq \max(\|x\|, \|y\|)$
- (3)  $\|x\| = 0 \Leftrightarrow x = 0$ .

We can easily prove that  $\|1\| = \|-1\| = 1$ , and that equality holds in equation (2) unless  $\|x\| = \|y\|$ .

As an example let  $K$  be the field of rational numbers. Any  $x \neq 0$  in  $K$  may be written as  $2^t(r/s)$ , where  $r$  and  $s$  are odd integers and  $t$  is an integer. Set  $\|x\| = (1/2)^t$ . In this way we get an ultranorm on the rationals in which  $\|2\| < 1$ . This ultranorm is known as the *2-adic absolute value*. The more general fact that we shall need is this: *There is an ultranorm on the field of real numbers (or more generally on any extension of the rational numbers) such that  $\|2\| < 1$ .* This follows from the theorem of the extension of valuations whose proof may be found in many places; for example see [3].

Granting the above facts we may argue as follows. Choose an ultranorm on the reals for which  $\|2\| < 1$ . Divide the points of the plane into three sets in the following way:

- (1)  $(x, y)$  is in  $\mathfrak{A}$  if  $\|x\| < 1$  and  $\|y\| < 1$ ,
- (2)  $(x, y)$  is in  $\mathfrak{B}$  if  $\|x\| \geq 1$  and  $\|x\| \geq \|y\|$ ,
- (3)  $(x, y)$  is in  $\mathfrak{C}$  if  $\|y\| \geq 1$  and  $\|y\| > \|x\|$ .

Suppose now that  $P = (x, y)$  and  $P' = (x', y')$  are points and that  $P'$  is a translate of  $P$  by a point of type  $\mathfrak{A}$ ; in other words, that both  $\|x' - x\| < 1$  and  $\|y' - y\| < 1$ . Then  $P$  and  $P'$  have the same type. If  $P$  is of type  $\mathfrak{A}$  this is obvious. If  $P$  is of type  $\mathfrak{B}$ , then  $\|x'\| = \|x\| \geq 1$ , while  $\|y'\| \leq \max(1, \|y\|) \leq \|x\| = \|x'\|$ ; so  $P'$  is of type  $\mathfrak{B}$  too. If  $P$  is of type  $\mathfrak{C}$  the argument is similar.

It is now easy to see that a line  $L$  cannot contain points of all three types. For by translating a point of type  $\mathfrak{A}$  on  $L$  to the origin we may assume that  $(0, 0)$  is on  $L$ . Let  $(x, y)$  and  $(x', y')$  be points of  $L$  of types  $\mathfrak{B}$  and  $\mathfrak{C}$ . Then  $\|x\| \geq \|y\|$ ,  $\|y'\| > \|x'\|$ , and  $\|xy'\| > \|x'y\|$ . This is absurd as  $xy' = x'y$ . Note also that if a triangle  $T$  has vertices of all three types then  $\|\text{area } T\| > 1$ . For we may assume that the vertex of  $T$  of type  $\mathfrak{A}$  is  $(0, 0)$ . Let  $(x, y)$  and  $(x', y')$  be the vertices of types  $\mathfrak{B}$  and  $\mathfrak{C}$ . Then  $\text{area } T$ , up to sign, is equal to  $\frac{1}{2}(xy' - x'y)$ . But  $\|xy'\| > \|x'y\|$ . So  $\|\text{area } T\| = \|\frac{1}{2}\| \|xy'\| = \|\frac{1}{2}\| \cdot \|x\| \cdot \|y'\| > 1$ .

Suppose now that  $S = [0, 1] \times [0, 1]$  is divided into  $m$  nonoverlapping triangles  $T_i$  each of area  $1/m$ . Obviously  $S$  has exactly one face of type  $\mathfrak{A}\mathfrak{B}$ ; by the lemma, some  $T_i$  has vertices of all three types. By the paragraph above,  $\|\text{area } T_i\| = \|1/m\| > 1$ . So  $m$  is even. (Note that if all vertices have rational coordinates, then we can argue directly with the 2-adic absolute value of the rationals, and avoid the theorem of the extension of valuations; this is essentially what was done in [2].)

Finally, we indicate the proof of the more general theorem mentioned at the beginning of this paper. Let  $A$  be the ring  $\mathbf{Z}[a_1, \dots, a_m]$ . If 2 generates the unit ideal in  $A$ , then  $1 = 2f(a_1, \dots, a_m)$  and we are done. If  $2A \neq A$ , then 2 is contained in a height 1 prime ideal  $P$  of  $A$ . The integral closure of the local ring of  $P$  on  $A$  is a discrete valuation ring. This ring gives rise to an ultranorm on the quotient field of  $A$  such that  $\|a_i\| \leq 1$ , while  $\|2\| < 1$ . Extend this ultranorm to the reals, and use it to subdivide the plane into points of three types as above. Then, as above, some  $T_i$  has vertices of all three types, and  $\|a_i\| = \|\text{area } T_i\| > 1$ ,

a contradiction. (By using valuation rings instead of ultranorms we could simplify the proof a little.)

The above proof is not so wildly nonconstructive as it first appears. For the entire argument is carried out in the field generated by the coordinates of the vertices. So it is only necessary to extend our ultranorm from  $Q$  to this finitely generated field, not to the entire field of real numbers.

#### References

1. Fred Richman and John Thomas, Problem 5479, this MONTHLY, 74 (1967) 329.
2. John Thomas, A dissection problem, Math. Mag., 41 (1968) 187-190.
3. Serge Lang, Algebra, Addison-Wesley, Reading, Mass., 1965, p. 299.

#### SYLVESTER'S PROBLEM ON COLLINEAR POINTS AND A RELATIVE

G. D. CHAKERIAN, University of California, Davis

In 1893, J. J. Sylvester posed the following question in the Educational Times: *Given a finite set of points in the plane, not all lying on a straight line, must there be a line containing exactly two of the points?*

This question was not settled until after 1930, when T. Gallai proved the affirmative answer. Many proofs have been given since that time, some sparked by a restatement of the problem by P. Erdős [3] in this MONTHLY. V. C. Williams [9] recently gave a proof in this MONTHLY. Extensive references to the literature which has grown up around the problem can be found in [2], [4], and [5].

If we settle upon the projective plane as an appropriate setting and state the problem in its dual form, we obtain: *Given  $n$  lines in the projective plane, not all concurrent, must there be a point lying on exactly two of the lines?*

Thinking of the projective plane as a sphere with antipodes identified, and then immediately observing that the identification contributes nothing to the problem, we obtain the following equivalent formulation of Sylvester's problem: *Given  $n$  great circles on a sphere, not all concurrent, must there be a point lying on exactly two of the great circles?*

In his exquisite paper on zonohedra, H. S. M. Coxeter [1] observes in passing that in this context the positive answer to Sylvester's question follows immediately from the fact that there exists no map on the sphere (each of whose countries has at least 3 sides) such that all vertices have valence 6 or more. For a set of  $n$  great circles such that no point lies on exactly two circles, would form a map on the sphere such that each vertex has valence at least 6.

Here is a short proof of this fact about maps. Suppose we are given a map on the sphere such that each face (country) has 3 or more sides. Suppose each vertex had valence at least 6 (the valence of a vertex is the number of edges emanating from that vertex). If  $V$  denotes the total number of vertices,  $E$  the number of edges, and  $F$  the number of faces, then we have Euler's relation,

$$(1) \quad V - E + F = 2.$$